



Provis Rudd Clinics

123 Edward Street Suite 400
Toronto, ON M5G 1E2

Tel: (416) 597-0997 | Fax: (416) 597-2912

PEGLYTE INSTRUCTIONS FOR COLONOSCOPY

Date&Time: _____ **Doctor:** _____

Purchase the following: 1 jug of Peglyte and 1 box of Dulcolax tablets, which can be purchased at our clinic

- You must arrange for someone to accompany you to and from the clinic as you will not be able to drive for 24 hours following sedation. The person picking you up must meet you in the recovery area of the clinic.
- Bring juice, Gatorade or Powerade for after your procedure to restore energy.
- Wear loose fitted clothing to your appointment and do not bring any valuables.
- If you require a translator they must be present with you for your appointment.
- No cannabis use 48 hours prior to your procedure.

4 days before your appointment: Stop eating all seeds, nuts and grains including multigrain breads, flax seeds, fruits/vegetables with seeds (ie: tomatoes, cucumbers, watermelon, all types of berries)

Day before your appointment: From the time you wake up you must remain on clear fluids only, no food. Clear fluids such as water, apple juice, clear or white Gatorade, 7-up, Sprite, ginger ale, white grape juice and white cranberry juice. No milk or dark fluids.

Appointment before 12pm:	Appointment 12pm or later:
<u>Day before your appointment:</u> At 1pm take 4 Dulcolax tablets with water Between 5:30pm-8:30pm drink 3 litres of Peglyte	<u>Day before your appointment:</u> At 1pm take 4 Dulcolax tablets with water Between 6pm-8pm drink 2 litres of Peglyte
<u>Morning of your appointment:</u> Between 5am-6am drink the 4 th litre of Peglyte	<u>Morning of your appointment:</u> Between 7am-9am finish the 2 litres of Peglyte
Remain on clear fluid only no food	

For colonoscopy: Stop drinking all fluids including water 2 hours before your appointment

Colonoscopy with Gastroscopy: Stop drinking all fluids including water 3 hours before your appointment

Medication Restrictions (please confirm with your family doctor):

Stop Xarelto, Pradaxa, Eliquis and Brilinta 2 days prior to procedure

Stop Coumadin, Warfarin and Plavix 5 days prior to procedure

Stop Ticlid and Iron supplements 7 days prior to procedure

Stop oral diabetic medications the evening before and the morning of the procedure. If you are on Insulin please discuss with your doctor about cutting the morning dose in half. Stop Ozempic 14 days before.

Medication for blood pressure, thyroid, cholesterol can be taken morning of appointment with a sip of water

You can take Tylenol or prescribed medication for pain control